



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MECHANICS.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

28. Proposed by O. W. ANTHONY, Professor of Mathematics, New Windsor College, New Windsor, Maryland.


A movable finite wire carrying a current of electricity is perpendicular to and on one side of an infinite wire also carrying a current. Investigate the motion of the movable wire.

Solution by the PROPOSER.

Let AB be the finite wire, and DC the infinite wire. Let the current flow away from the infinite wire in the short one. Also call μ_1 , μ_2 the current strengths of the two currents, and m the power of the

finite wire. Then $df_{PQ} = \mu_1 \mu_2 \frac{xdx dz}{x^2 + z^2}$, $x = QS$, $z = PS$. Resolving forces perpendicular and parallel to DC we have

$$\begin{aligned} df_{PR} &= \mu_1 \mu_2 \frac{xdx dz}{(x^2 + z^2)^{\frac{3}{2}}}. \\ \therefore f_{PR} &= 2\mu_1 \mu_2 \int_{z_1}^{z_2} \int_0^\infty \frac{xdx dz}{(x^2 + z^2)^{\frac{3}{2}}}. \\ &= 2\mu_1 \mu_2 \log\left(\frac{z_2}{z_1}\right). \\ \therefore \frac{d^2 s}{dt^2} &= \frac{2}{m} \mu_1 \mu_2 \log\left(\frac{z_2}{z_1}\right). \\ s &= \frac{1}{m} \mu_1 \mu_2 \log\left(\frac{z_2}{z_1}\right) t^2 + k_1 t + k_2. \end{aligned}$$



29. Proposed by J. A. CALDERHEAD, A. B., Superintendent of Schools, Limaville, Ohio.

Show that if a body be projected from the angle A of a plane triangle ABC so as to strike the side CB at a point D , then, if its course after reflection at D be parallel to AB ,

$$\tan \angle DAB = \frac{(1+e)\cot B}{1-e\cot^2 B}.$$

Solution by ALFRED HUME, C. E., D. Sc., Professor of Mathematics in the University of Mississippi, University P. O., Mississippi.

The angle between the course of the body before impact and the side CB is $180^\circ - (B + DAB)$.

$$\therefore e = \frac{\tan B}{\tan[180^\circ - (B + DAB)]};$$